

Original Research Paper

Investigating the energy consumption and carbon footprint in mung bean production ecosystems in Iran

Amin Fathi^{1,2}, Kamran Kheiralipour^{1*}¹Mechanical Engineering of Biosystems Department, Ilam University, Ilam, Iran²Department of Agronomy, Ayatollah Amoli Branch, Islamic Azad University, Amol, Iran

Biosystems Engineering and Renewable Energies 2025, 1 (1): 59-63

KEYWORDS

Mung bean
Input-output
Energy
Carbon dioxide
Environmental indicators

* Corresponding author
k.kheiralipour@ilam.ac.ir

Article history

Received: 2024-10-21

Revised: 2024-12-28

Accepted: 2024-12-30

ABSTRACT

Energy consumption is one of the main issues in production systems because of its role in production costs and environmental impacts. This investigation was conducted to assess the environmental aspect of mung bean farms in the 2022-2023 crop year. The data from 78 mung bean farms in Darreh Shahr, Ilam, Iran were collected. Input-output materials were obtained and energy indicators and carbon footprint were calculated based on them. The results showed that the carbon footprint of mung bean production was 2.74 ton CO₂ eq. ha⁻¹ (2.42 kg CO₂ eq. kg⁻¹). The total amount of input and output energy in mung bean production was 12586.49 and 16604.40 MJ ha⁻¹, respectively. The most important energy-intensive inputs in mung bean production were nitrogen fertilizer, fuel, and electricity with the shares of 32.57, 25.88, and 24.43%, respectively. The value of energy proportion was 1.32. Energy productivity and net energy gain efficiency had the values of 0.09 kg MJ⁻¹ and 0.32 MJ MJ⁻¹, respectively. Given the low share of renewable energy (3.49%), increasing renewable energy consumption is necessary for mung bean production.

1. Introduction

Legumes are beneficial in terms of human diet, animal nutrition, and soil fertility. One of the members of the legume family is the mung bean (*Vigna Radiata* L.) which is an annual, short-day, thermophilic, short-growing, and summer plant cultivated in tropical and subtropical regions. Also, this crop is a rich source of vegetable protein (Gilani et al., 2021). Like most legume families, mung bean has a high nitrogen absorption capacity and is therefore considered an important crop in agriculture (Hunady and Hochman, 2014; Kazemi et al., 2016). Mung bean has a worldwide cultivation area of 7.3×10⁶ ha with an annual yield of 5.3×10⁶ ton (Nair and Schreinemachers, 2020). Its farm area is 25,000 ha in Iran (FAOESTAT, 2023).

Sustainability is a main issue in production systems due to the limited resources (Kheiralipour, 2022). Energy and carbon footprint are the key elements in the sustainable production path. With the increasing world population and resource limitations, access to sufficient energy will be more difficult in the future (Khodaei et al., 2022). Currently, ensuring food security for the growing world population by preserving basic land and water resources with minimal environmental impacts has become one of the fundamental challenges in sustainable agriculture (Mohammadzadeh et al., 2017). The limited energy sources and the impacts of the misuse of non-renewable energy sources such as fossil fuels on the environment and human health are certain which emphasizes the study of energy consumption patterns and energy-intensive inputs in ecosystems (Snyder et al., 2009; Kheiralipour et al., 2017, 2018).

Energy is the ability to create change in the form of work or heat. The energy status is studied in different production systems to decrease energy consumption and increase energy

productivity. It is basically consumed in two different forms: renewable and nonrenewable. The goal of sustainable production is to increase the share of renewable energy sources and decrease the share of nonrenewable energy sources to conserve the environment.

In agriculture, like other sectors, energy is directly consumed to conduct different operations such as soil preparation (leveling and plowing), planting, protecting, harvesting, and postharvest processing. Also, it is indirectly consumed in the production of inputs in production units such as pesticides, chemical fertilizers, machinery, fuel, and seed (Kheiralipour and Sheikhi, 2020; Fathi et al., 2020; Khodaei et al., 2022). Energy consumption in agricultural ecosystems has led to increased production productivity and economic growth. However, the development of industrial agriculture, especially intensive agricultural systems that are highly dependent on fertilizers, chemical pesticides, and other energy-intensive inputs has increased greenhouse gas emissions (Li et al., 2016).

On a global scale, about 5% of total energy consumption is related to the agricultural sector, but about 11% of greenhouse gas emissions are related to this sector (Smith et al., 2014). These emissions are mainly due to the consumption of fossil fuels, pesticides, chemical fertilizers, electricity, and conducting tillage operations (Camargo et al., 2013). Also, energy efficiency and the ratio of greenhouse gas emissions to the outputs in agricultural ecosystems can be affected by different factors such as the type of cropping system, cropping pattern, applied technology, population employed in agriculture, farmer knowledge, type and amount of used chemical fertilizer, and crop yield (Fathi et al., 2020; Mohammadzadeh et al., 2017). Therefore, assessing the efficiency of input use in agricultural ecosystems plays a significant role in reducing energy consumption and production costs and also in designing sustainable and environmentally

friendly ecosystems (Payandeh et al., 2017; Kheiralipour, 2020; Dekamin et al., 2022; Pourmehdi and Kheiralipour, 2023).

Energy consumption of various agronomy (Molaei et al., 2008; Pourmehdi and Kheiralipour, 2024; Ramedani et al., 2024), horticultural (Hesampour et al., 2022; Dekamin and Kheiralipour, 2023), and animal (Payandeh et al., 2016; Ramedani et al., 2019; Heidarbeigi and Sheikh, 2023) products in agriculture and food products (Kheiralipour et al., 2018; Jalilian et al., 2020; Gholamrezaee et al., 2021) have been studied in the literature. In terms of carbon footprint, Heidari et al. (2017) evaluated wheat systems in Iran. Zhang et al. (2017) calculated the carbon footprint of maize, wheat, and rice in China. Kashya and Agarwal (2021) studied rice and wheat systems in Punjab. However, Abad-Gonzalez et al. (2024) investigated the energy status of mung bean in Iran, but they did not report its energy indicators and carbon footprint. Given the importance of mung bean, it is necessary to examine how to produce this crop sustainably. Therefore, the present study aimed to investigate the status of energy and carbon footprint in mung bean farms.

2. Materials and Methods

Figure 1 depicts the steps in the present research to determine the energy status and carbon footprint of mung bean production in Darreh Shahr, Ilam, Iran.

2.1. Data collection

A questionnaire was designed to gather information on mung bean systems. Interviews with 78 ght mung bean farm owners were done in the 2022-2023 crop year in Darreh Shahr Township, Ilam, Iran. Darreh Shahr is located 135 km southeast of Ilam and 160 km. The climate of the region is temperate and mountainous, and its 30-year average rainfall is 413 mm.

2.2. Energy calculations

To calculate the energy input, the amount of materials inputs in mung bean production including human labor, machinery, electricity, diesel fuel, seeds, fertilizers, and chemical sprays per hectare was determined. Then, they were multiplied by the energy equivalent corresponding to each of them (Table 1). The contribution of each input to the total energy input was calculated, as well as the values of different energy forms. Energy indicators were calculated using the following equations (Pourmehdi and Kheiralipour, 2024)

$$ER = \frac{OE}{IE} \times 100 \quad (1)$$

$$EP = \frac{OY}{IE} \quad (2)$$

$$EI = \frac{IE}{OY} \quad (3)$$

$$NEG = OE - IE \quad (4)$$

$$NEGE = \frac{OE - IE}{IE} \quad (5)$$

where ER is the efficiency of energy consumption, EP is the productivity of energy consumption (kg MJ⁻¹), EI is the energy intensity (MJ kg⁻¹), NEG is the net energy gain (MJ ha⁻¹), NEGE is the net energy gain efficiency indicator, OE is the total output energy (MJ), IE is the total input energy (MJ), and OY is the amount of output materials (kg).

2.3. Carbon footprint calculation

Greenhouse gases (GPW) are emitted from various activities. The main GPWs in agricultural ecosystems are carbon dioxide, methane, and nitrous oxide. The effect of each of carbon dioxide, methane, and nitrous oxide on global warming is different, such that methane and nitrous oxide contribute to global warming about 21 and 310 times more than carbon dioxide, respectively

(IPCC, 2007). Therefore, the amounts of these emissions are calculated as carbon dioxide equivalent. The amounts of greenhouse gases were calculated using the corresponding emission factors in Table 2. The global warming potential was calculated based on Eq. (6) (Kramer et al., 1999)

$$GWP = CO_2\text{flux} + (N_2O\text{flux} \times 310) + (CH_4\text{flux} \times 21) \quad (6)$$

where GWP, CO₂flux, N₂flux, and CH₄flux represent the global warming potential (kg CO₂ eq. ha⁻¹), carbon dioxide, nitrous oxide, and methane emission, respectively.

3. Results and Discussion

3.1. Energy

The average values of input and output energy in mung bean production are shown in Table 3. The total yield in mung bean production was 1129.55 kg ha⁻¹ and its energy equivalent was 16604.40 MJ ha⁻¹. The total energy of inputs was equal to 12586.49 MJ ha⁻¹ (11142.92 MJ ton⁻¹). This amount was lower than that of mung bean production in Golestan Province, Iran (27400 MJ ton⁻¹) (Abad-Gonzalez et al., 2024). The average energy input in corn in Iran has been reported to be 53602 MJ ha⁻¹ (Beheshti Tabar et al., 2010). Pourmehdi and Kheiralipour (2024) reported that the total energy input consumed in rainfed and irrigated wheat in Cherdavol township, Ilam, Iran, was 12085 and 20812 MJ ha⁻¹, respectively, and the total output (grain and straw) energy was 61933 and 132492 MJ ha⁻¹, respectively.

Trudy et al. (2018) stated that among all inputs, energy consumption related to nitrogen ranked first with 38.03% in wheat production followed by fuel and seed. In a research conducted to investigate the energy status of faba beans in Iran, it was found that the main energy contributor was nitrogen fertilizer (Kazemi et al., 2015). Pourmehdi and Kheiralipour (2024) reported that diesel fuel (79.6 l ha⁻¹) was the main contributor to dryland wheat (37.13%) followed by nitrogen fertilizer (35.51%) and seeds (18.24%). The main energy contributors in wheat production in irrigated systems were electricity, fuel, nitrogen, and seeds with the shares of 29, 27, 20, and 15%, respectively.

The values of energy efficiency, energy intensively, energy productivity, net energy gain, and net energy gain efficiency are given in Table 4. Energy efficiency in mung bean cultivation was calculated to be 1.32. The energy efficiency and net energy gain in mung bean production were 0.09 kg MJ⁻¹ and 4017.91 MJ ha⁻¹, respectively. Kazemi et al. (2015) reported that the energy efficiency in faba bean production was 4.70. Energy productivity, net energy gain, and energy intensity had the values of 0.23 kg MJ⁻¹, 51226.30 MJ ha⁻¹, and 4.25 MJ kg⁻¹, respectively. According to the results of Pourmehdi and Kheiralipour (2024), the energy ratios of wheat production in irrigated and dryland systems were 37.6 and 12.5, respectively. The values of the energy efficiency in the fields were 0.38 and 0.48, respectively. The net energy gain efficiency in the farms had values of 5.37 and 4.12, respectively.





Figure 1. The steps conducted in the present research.

Table 1. The energy equivalent of input-output material in mung bean production.

Material (reference)	Unit	Equivalent energy (MJ Unit ⁻¹)
Input		
Diesel fuel (De et al., 2001)	l	56.31
Oil (Kitani, 1999)	l	47.80
Labor (Singh et al., 2002)	h	1.96
Machinery (Chauhan et al., 2006)	kg	62.70
Pesticide (Ozkan et al., 2004)	kg	199.00
Herbicide (Erdal et al., 2007)	kg	238.00
Nitrogen (Esengun et al., 2007))	kg	66.14
Phosphorous (Esengun et al., 2007))	Kg	12.44
Potassium (Esengun et al., 2007))	kg	11.15
Electricity (Ozkan et al., 2004))	kWh	11.93
Seed (Yousefi et al., 2014b)	kg	14.7
Output		
Mung bean (Yousefi et al., 2014a))	kg	14.7

Table 2. The greenhouse gases (g Unit⁻¹) of different input materials in mung bean production (Fathi et al., 2020; Yousefi et al., 2014a,b).

Inputs (Unit)	CO ₂	N ₂ O	CH ₄
Diesel (l)	3560.00	0.70	5.20
Nitrogen fertilizer (kg)	3100.00	0.03	3.70
Phosphate fertilizer (P ₂ O ₅) (kg)	1000.00	0.02	1.80
Potassium fertilizer (K ₂ O) (kg)	700.00	0.01	1.00
Electricity (kWh)	61.20	8.82	0.02



**Figure 2.** The energy contributors in mung bean production.**Table 3.** Material and energy amounts and the shares of inputs in mung bean production.

Material	Amount (Unit ha ⁻¹)	Energy (MJ ha ⁻¹)
Input		
Labor (h)	73.09	143.26
Machinery (kg)	16.15	1012.85
Diesel fuel (l)	63.45	3256.82
Oil (l)	1.14	54.49
Fertilizer (kg)		
Nitrogen	61.99	4099.83
Phosphorous	3.33	41.47
Potassium	3.33	37.17
Sprays (l)		
Pesticide	0.65	154.96
Herbicide	1.45	415.78
Electricity (kWh)	835.99	3074.35
Water (m ³)	4134.16	-
Seed (kg)	20.10	295.51
Total	-	12586.49
Output		
Mung bean	1129.55	16604.40
Total	-	16604.40

Table 4. Energy indicators in mung bean production.

Indicator	Value
Energy efficiency	1.32
Energy productivity	0.09
Energy intensity	11.11
Net energy gain	4017.91
Net energy gain efficiency	0.32

As Figure 3 shows, the difference between the shares of direct and indirect energy was low (2%). The share of direct and indirect energy in mung bean production was 51.44 and 48.56%, respectively. In mung bean production, the direct energy included labor, diesel, and electricity, and the indirect energy included seed, machinery, fertilizer, spray, and oil. The renewable energy sources had been consumed very low (3.49%) compared to the nonrenewable sources (96.51%) (Figure 4). The nonrenewable energy sources were diesel, oil, electricity, machinery, fertilizers, and sprays, and renewable energy sources were labor and seed.

**Figure 3.** Direct and indirect energy forms.**Figure 4.** The shares of renewable and nonrenewable energy.

In faba bean farms, Kazemi et al. (2015) showed that indirect energy (81.23%) and non-renewable energy (74.72%) sources were more consumed than the direct (18.77%) and renewable energy forms (25.28%), respectively. Pourmehdi and Kheiralipour (2024) estimated that the share of direct and indirect energy of the total energy consumed for the production of one hectare of dryland wheat in Chardavol township was 38.05 and 61.95%, respectively. Also, the share of direct (57.14%) of the total energy consumed for the production of one hectare of wheat farms with the irrigated system was higher than that of indirect energy (42.86%).

3.2. Carbon footprint

The values of carbon footprint in mung bean cultivation are shown in Table 5. As reported in this table, the carbon footprint of mung bean was 2737.24 kg CO₂ eq. ha⁻¹ (2.74 ton CO₂ eq. ha⁻¹). Since 1.13 ton of mung bean was produced in each hectare (Table 5), the carbon footprint was 2.42 kg CO₂ eq. kg⁻¹. This value is higher than the carbon footprint of wheat, maize, and rice. Heidari et al., (2017) reported 1.60 kg CO₂ eq. kg⁻¹ as the mean of carbon footprint (0.80 to 3.00 kg CO₂ eq. kg⁻¹) of durum wheat production in Iran. Zhang et al. (2017) reported the carbon footprint of maize, wheat, and rice as 4052, 5455, and 11881 kg CO₂ eq. ha⁻¹ respectively in China (0.48, 0.75, and 1.60 kg CO₂ eq. kg⁻¹, respectively). Kashya and Agarwal (2021) reported that the mean values of the carbon footprint of rice and wheat systems in different regions in Punjab, India, were 1.20±0.70 and 0.83±0.23 ton CO₂ eq. ton⁻¹, respectively.

The shares of the inputs in the carbon footprint of mung bean production are shown in Figure 5. The main contributors to carbon footprint were 87.23, 9.01, and 3.32% belonged to electricity, diesel fuel, and nitrogen fertilizer with the shares of 2387.59 kg CO₂ eq. ha⁻¹, 246.57 kg CO₂ eq. ha⁻¹, and 90.87 kg CO₂ eq. ha⁻¹.

4. Conclusions

The energy indicators and carbon footprint of mung bean farms in Darreh Shahr, Ilam, Iran, were investigated in the present research. The most important energy-intensive inputs in mung bean production were nitrogen fertilizer, fuel, and electricity with the shares of 32.57, 25.88, and 24.43%, respectively. The most important contributors to the carbon footprint of mung bean production were electricity, diesel fuel, and nitrogen fertilizer

References

- Abad-Gonzalez, J., Nadi, F., Perez-Neira, D. 2024. Energy-water-food security nexus in mung bean production in Iran: An LCA approach. *Ecological Indicators*, 158, 111442.
- Beheshti Tabar, I., Keyhani, A.R. and Rafiee, S. 2010. Energy balance in Iran Agronomy. *Renew. Energy*. 14: 489- 855.
- Camargo, G. G., Ryan, M. R., & Richard, T. L. (2013). Energy use and greenhouse gas emissions from crop production using the farm energy analysis tool. *BioScience*, 63(4), 263-273.
- Chauhan, N.S., Mohapatra, P.K.J., Pandey, K.P. (2006) Improving energy productivity in paddy production through benchmarking: an application of data envelopment analysis. *Energy Convers Manag*, 47, 1063-85.
- De, D., Singh, R.S., Chandra, H. (2001) Technological impact on energy consumption in rainfed soybean cultivation in Madhya Pradesh. *Appl Energy*, 70, 193-213.
- Dekamin, M., Kheiralipour, K., Keshavarz Afshar, R., 2022. Energy, economic, and environmental assessment of coriander seed production using material flow cost accounting and life cycle assessment. *Environmental Science and Pollution Research*.
- Dekamin, M., Kheiralipour, K., 2023. Material and Energy Flow Cost Accounting (MEFCA) of Grape Production in Malayer City. *Iranian Agricultural Economics Society*. 37(3), 325-340.
- Erdal, G., Esengu, K., Erdal, H., Gunduz, O. (2007) Energy use and economical analysis of sugar beet production in Tokat province of Turkey. *Energy*, 32, 35-41.
- Esengun, K., Erdal, G., Gündüz, O., & Erdal, H. (2007). An economic analysis and energy use in stake-tomato production in Tokat province of Turkey. *Renewable Energy*, 32(11), 1873-1881.

with the shares of 87.23, 9.01, and 3.32%, respectively. Using renewable energy sources, bio-fertilizers, and minimum tillage equipment can decrease the input energy and carbon footprint of mung bean production.

Table 5. The carbon footprint (kg CO₂ eq. ha⁻¹) in mung bean production.

Inputs	CO ₂	N ₂ O	CH ₄	Carbon footprint
Diesel	225.88	0.04	0.33	246.57
Nitrogen	88.39	1.00×10 ⁻³	0.11	90.87
Phosphorous	1.53	0.00	0.00	1.60
Potassium	0.79	0.00	0.00	0.82
Herbicide	7.30	-	-	7.30
Insecticide	2.47	-	-	2.47
Electricity	52.26	7.53	0.02	2387.59
Total	378.63	7.58	0.46	2737.24
CO ₂ eq.	378.63	2349.02	9.58	2737.24




Figure 5. The share of different inputs in the carbon footprint of mung bean production.

Acknowledgments

The authors thank Ilam University for supporting this investigation. Also, the mung bean farmers in Darreh Shahr, Ilam, Iran, who provided the information used in this research are appreciated.

- FAOSTAT (2023). Food and Agriculture Organization of the United Nations (FAO). FAOSTAT Database. Online: <https://www.fao.org/faostat/en/#data>.
- Fathi, A., Barari Tari, D., Fallah Amoli, H., & Niknejad, Y. (2020). Study of energy consumption and greenhouse gas (GHG) emissions in corn production systems: influence of different tillage systems and use of fertilizer. *Communications in Soil Science and Plant Analysis*, 51(6), 769-778.
- Gholamrezaee, H., Kamran Kheiralipour, K., Rafiee, S., 2021. Investigation of energy and environmental indicators in sugar production from sugar beet. *Journal of Environmental Studies Sciences* 6(2): 3540-3548.
- Ghorbani, R., Mondani, F., Amirmoradi, S., Feizi, H., Khorramdel, S., Teimouri, M., ... & Aghel, H. (2011). A case study of energy use and economical analysis of irrigated and dryland wheat production systems. *Applied Energy*, 88(1), 283-288.
- Gilani, A., Abbasdokht, H., & Gholami, A. (2021). Nutritional effects of essential sulfur along with *Halothiobacillus neapolitanus* on morphophysiological and agronomic responses of mung bean (*Vigna radiata* L.). *JOURNAL OF AGRICULTURAL SCIENCE AND SUSTAINABLE PRODUCTION*, 31(4), 189-205.
- Guinee, J.B. 2002. Handbook on life cycle assessment operational guide to the ISO standards. NewYork: Kluwer Academic.
- Heidarbeigi, K., Sheikhi, N. 2023. Investigating the energy consumption indicators and environmental effect of the trout ponds in Ilam province. *Journal of Agricultural Mechanization*, 8(1), 15-22.
- Heidari, M.D., Mobli, H., Omid, M., Rafiee, S., Jamali Marbini, V., Elshout, P.M.F., Van Zelm, R., Huijbregts, M.A.J. 2017. Spatial and technological variability

- in the carbon footprint of durum wheat production in Iran. *International Journal of Life Cycle Assessment*. DOI 10.1007/s11367-017-1283-1.
- Hesampour, R., Hassani, M., Hanafiah, M.M., Heidarbeigi, K. 2022. Technical efficiency, sensitivity analysis and economic assessment applying data envelopment analysis approach: A case study of date production in Khuzestan State of Iran. *Journal of the Saudi Society of Agricultural Sciences*. 21(3), 197-207.
- Hunady, I., Hochman, M., 2014. Potential of legume-cereal intercropping for increasing yields and yield stability for self-sufficiency with animal fodder in organic farming. *Czech Journal of Genetics and Plant Breeding* 50, 185–194.
- IPCC, 2006. IPCC 14040. IPCC guidelines for national greenhouse gas inventories. In: Eggleston HS, Buendia L, Miwa K, Ngara T, Tanabe K, editors. Prepared by the national greenhouse gas inventories programme. Japan: IGES.
- ISO 14040, 2006; Environmental Management–Life Cycle Assessment–Principles and Framework. International Organization for Standardization: Geneva, Switzerland, 2006.
- Jalilian, M.M., Kheiralipour, K., Mirzaee Ghaleh, E. 2021. Comparison of environmental indicators in Sangak and Lavash bread production in Eslamabad-e-Gharb, Kermanshah. *Journal of Environmental Sciences Studies* 5(4), 3198-3203.
- Jalilian, M.M., Kheiralipour, K., Mirzaee Ghaleh, E. 2020. Comparison of energy status in the production process of Lavash and Sangak bread. 12th National Iranian Congress on Mechanical Engineering of Biosystems and Mechanization. 3-5 February, Ahvaz, Iran.
- Kashyap, D., Agarwal, T. 2021. Carbon footprint and water footprint of rice and wheat production in Punjab, India. *Agricultural Systems*, 186, 102959.
- Kazemi, H., Shahbyki, M., Baghbani, S., 2015. Energy analysis for faba bean production: A case study in Golestan province. *Iran. Sustainable Production and Consumption* 3, 15-20.
- Kheiralipour, K. (2022) *Sustainable Production: Definitions, Aspects, and Elements*. 1st Edition. Nova Science Publishers, New York, US.
- Kheiralipour, K. 2021, *Environmental Life Cycle Assessment of Poultry Production Systems*. In: Jacob-Lopes, E., Zepka, L. Q., Depra, M. C. Interdisciplinary applications of the life cycle assessment tool. 1st Ed., Nova Science Publishers, New York, USA.
- Kheiralipour, K. 2020. *Environmental Life Cycle Assessment*. 1st Ed., Ilam University Publication. Ilam, Iran.
- Kheiralipour, K., Brandao, M., Holka, M., Chorynski, A. 2024a. A review of environmental impacts of wheat production in different agrotechnical systems. *Resources*, 13(7), 93.
- Kheiralipour, K., Gholamrezaee, G., Rafiee, S. (2018) Investigation of energy status in sugar production. 11th National Congress on Biosystems Engineering and Agricultural Mechanization, 3-5 September, Hamedan, Iran.
- Kheiralipour, K., Jafari Samrin, H., Soleimani, M. 2017. Determining the environmental impacts of canola production by life cycle assessment, case study: Ardabil Province. *Iranian Journal of Biosystems Engineering*, 48(4), 517-526.
- Kheiralipour, K., Khoobbakht, M., Karimi, M. 2024. Effect of biodiesel on environmental impacts of diesel mechanical power generation by life cycle assessment. *Energy*, 289, 15, 129948.
- Kheiralipour, K., Payandeh, Z., Khoshnevisan, B. (2017). Evaluation of environmental impacts in turkey production system in Iran. *Iranian Journal of Applied Animal Science*, 7, 507-512.
- Kheiralipour, K., Rafiee, S., Karimi, M., Nadimi, M., Paliwal, J. 2024b. The environmental impacts of commercial poultry production systems using life cycle assessment: a review, *World's Poultry Science Journal*. 80 (1), 33-54.
- Kheiralipour, K., Sheikhi, N. (2021) Material and energy flow in different bread baking types. *Environment, Development and Sustainability*, 23, 10512-10527.
- Kheiralipour, K., Tashanifar, E., Hemati, A., Motaghd, S., Golmohammadi, A.R. 2022. Environmental impact investigation of natural gas refinery process based on LCA CML-IA baseline method. *Gas Processing Journal*. 9(2): 1-14.
- Khodaei johan, A., taki, M., & matoorian, H. (2022). Evaluating energy productivity, greenhouse gas emission, global warming potential and sustainability index of Wheat and Rapeseed agroecosystems in Khorramshahr. *Journal of Agricultural Science and Sustainable Production*, 32(1), 309-324.
- Kitani, O. (1999) *Energy and biomass engineering*. CIGR handbook of agricultural engineering. ASAE Publications, St Joseph, MI.
- Kramer KJ, Moll HC, Nonhebel S. 1999. Total greenhouse gas emissions related to the Dutch crop production system. *Agric Ecosyst Environ*. 72:9–16.
- Li, T., Baležentis, T., Makutėnienė, D., Streimikiene, D., & Kriščiukaitienė, I. (2016). Energy-related CO₂ emission in European Union agriculture: Driving forces and possibilities for reduction. *Applied Energy*, 180, 682-694.
- Mohammadi, A., & Omid, M. (2010). Economical analysis and relation between energy inputs and yield of greenhouse cucumber production in Iran. *Applied Energy*, 87(1), 191-196.
- Mohammadzadeh, A., Mahdavi Damghani, A., Vafabakhsh, J., & Dehimefard, R. (2017). Assessing energy efficiencies, economy, and global warming potential (GWP) effects of major crop production systems in Iran: a case study in East Azerbaijan province. *Environmental Science and Pollution Research*, 24(20), 16971-16984.
- Molaei, K., Keyhani, A., Karimi, M., Kheiralipour, K., Ghasemi Varnamkhasti, M. (2008) Energy ratio in dryland wheat-case study: Eghlid Township. *Iranian Journal of Biosystems Engineering*, 39(1), 13-19.
- Nair, R., Schreinemachers, P., 2020. Global Status and Economic Importance of Mungbean. In: Nair, R., Schafleitner, R., Lee, S.H. (Eds.), *The Mungbean Genome*. Compendium of Plant Genomes. Springer, Cham.
- Ozkan, B., Akcaoz, H., and Fert, C. (2004) Energy input-output analysis in Turkish agriculture. *Renewable Energy*, 29, 39-51.
- Payandeh, Z., Kheiralipour, K., Karimi, M. (2016) Evaluation of energy efficiency of broiler production farms using data envelopment analysis technique, case study: Isfahan Province. *Iranian Journal of Biosystems Engineering*, 47(3), 577-585. (In Persian).
- Payandeh, Z., Kheiralipour, K., Karimi, M., Khoshnevisan, B. (2017) Joint data envelopment analysis and life cycle assessment for environmental impact reduction in broiler production systems. *Energy*. 127, 768-774.
- Pourmehdi, K., Kheiralipour, K. 2024. Net energy gain efficiency, a new indicator to analyze energy systems, case study: Comparing wheat production systems. *Results in Engineering*, 22, 102211.
- Pourmehdi, K., Kheiralipour, K. 2023. Compression of input to total output index and environmental impacts of dryland and irrigated wheat production systems. *Ecological Indicators*, 148, 110048.
- Pourmehdi, K., Kheiralipour, K. 2020. Assessing the effects of wheat flour production on the environment. *Advances in Environmental Technology*. 2: 111-117.
- PRÉ 2006. PRÉ Consultants. SimaPro Database Manual.
- Ramedani, Z., Alimohammadian, L., Kheiralipour, K., Delpisheh, P., Abbasi, Z. 2019. Comparing energy state and environmental impacts in ostrich and chicken production systems. *Environmental Science and Pollution Research*. 27: 28284-28293.
- Ramedani, Z., Veisi, A., Lotfi, M., Kheiralipour, K., 2024. Comparison of material and energy indicators in sunflower and pumpkin seed production systems. *Sustainability Research, Technology Development and Assessment*. Doi.10.22098/SR.2024.14811.1020.
- Singh, H., Mishra, D., Nahar, N.M. (2002) Energy use pattern in production agriculture of a typical village in Arid Zone India Part I. *Energy Convers Manage*, 43(16), 2275-86.
- Smith, P., Bustamante, M., Ahammad, H., Clark, H., Dong, H., Elsidig, E., Haberl, H., Harper, R., House, J., and Jafari, M. 2014. *Agriculture, Forestry and Other Land Use (AFOLU)*. Mitigation of Climate Change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge: Cambridge University Press.
- Snyder, C. S., Bruulsema, T. W., Jensen, T. L., & Fixen, P. E. (2009). Review of greenhouse gas emissions from crop production systems and fertilizer management effects. *Agriculture, Ecosystems & Environment*, 133(3-4), 247-266.
- Yousefi, M., Khoramivafa, M., Mondani, F., 2014a. Integrated evaluation of Energy use, Greenhouse gas emissions and Global warming potential for sugar beet (*Beta vulgaris*) agroecosystems in Iran. *Atmos. Environ*. 92, 501-505.
- Yousefi, M., Mahdavi Damghani, A., Khoramivafa, M., 2014b. Energy consumption, greenhouse gas emissions and assessment of sustainability index in corn agroecosystems of Iran. *Sci Total Environ*. 493, 330-335.
- Zhang, D., Shen, J., Zhang, F., Li, Y., Zhang, W. 2017. Carbon footprint of grain production in China. *Scientific Reports*, 7, 4126.